



## CELL LAB Goat F(ab')<sub>2</sub> Anti-Human IgD ( $\delta$ chain specific)

<u>Cat. No.</u>	<u>Form</u>	<u>Quantity</u>
732573	Purified (UNLB) Antibody	0.5 mg
732574	Fluorescein (FITC) Conjugate	0.5 mg
732575	Biotin (BIOT) Conjugate	0.5 mg
732576	Phycoerythrin (PE) Conjugate	0.25 mg

**For Research Use Only. Not for use in diagnostic procedures.**

### DESCRIPTION

**Source:** Pepsin digest of goat anti-human IgD.  
**Cross Absorption:** Human IgM, IgG and IgA.  
**Purification:** Gel filtration chromatography of pepsin digested antibody.  
**Specificity:** Reacts with the heavy chain of human IgD as demonstrated by ELISA and flow cytometry. May also react with mouse IgD.

### APPLICATIONS

- Direct immunofluorescent staining of IgD<sup>+</sup> human B lymphocytes
- Enzyme-Linked Immunosorbent Assay (ELISA)
- Western blotting
- Dot- and slot-immunoblotting
- Immunohistochemistry

### CHARACTERIZATION

To ensure lot-to-lot consistency, each batch of product is tested to conform with characteristics of a standard reference reagent using ELISA, Particle Concentration Fluorescence Immunoassay and/or flow cytometry.

### WORKING DILUTIONS

**Immunofluorescence:** FITC conjugate  $\leq 1 \mu\text{g}/10^6$  cells  
BIOT conjugate  $\leq 1 \mu\text{g}/10^6$  cells  
PE conjugate  $\leq 0.1 \mu\text{g}/10^6$  cells

**ELISA:** BIOT conjugate 1:5,000-1:20,000

**Other Applications:** Since applications vary, determine the optimum working dilution of the product that is appropriate for your specific needs.

### HANDLING AND STORAGE

- The purified (UNLB) antibody is supplied as 0.5 mg of purified immunoglobulin in 1.0 mL of 100 mM borate buffered saline, pH 8.2. No preservatives or amine-containing buffer salts added.
- The fluorescein (FITC) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN<sub>3</sub>.
- The biotin (BIOT) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN<sub>3</sub>.

- The Phycoerythrin (PE) conjugate is supplied as 0.25 mg in 1.0 mL of PBS/NaN<sub>3</sub> and a stabilizing agent.
- Protect fluorochrome-conjugated forms from light. Do not freeze.
- Reagent is stable until the expiration date on the vial when stored at 2-8°C.

#### **STATEMENT OF WARNINGS**

1. Specimens, samples and all material coming in contact with them should be handled as if capable of transmitting infection and disposed of with proper precautions.
2. Never pipet by mouth and avoid contact of samples with skin and mucous membranes.
3. Do not use reagent beyond the expiration date on the vial label.
4. Minimize exposure of reagent to light during storage or incubation.
5. Avoid microbial contamination of reagent or erroneous results may occur.
6. Use Good Laboratory Practice (GLP) when handling this reagent.
7. Harmful if swallowed.
8. After contact with skin, wash immediately with plenty of water.
9. Contains sodium azide. Sodium azide under acidic conditions yields hydrazoic acid, an extremely toxic compound. Azide compounds should be flushed with running water while being discarded. These precautions are recommended to avoid deposits in metal piping in which explosive conditions can develop. If skin or eye contact occurs, immediately wash excessively with water.

#### **TRADEMARKS**

The Beckman Coulter logo is a trademark of Beckman Coulter, Inc.

For additional information or if damaged product is received, contact your local Beckman Coulter Representative.



Manufactured for:  
Beckman Coulter, Inc.  
4300 N. Harbor Blvd.  
Fullerton, CA 92835  
[www.beckmancoulter.com](http://www.beckmancoulter.com)

Printed in USA  
Made in USA

© 2005 Beckman Coulter, Inc.  
All Rights Reserved.

PN 734002-A