

Monoclonal Antibody IOTest® CD22-PC7

PN A74788 – 100 tests – Liquid – 10 µL/test* – Clone SJ10.1H11

Analyte Specific Reagent.

Analytical and performance characteristics are not established.

SPECIFICITY

The CD22 molecule is a single chain, type I transmembrane molecule, with a molecular weight of 130–140 kDa, composed by seven immunoglobulin-like (Ig-like) domains (1). CD22 is, like CD33 and the myelin-associated glycoprotein (MAG), a member of the sialoadhesin family (2). The N-terminal domain distal to the membrane is a V-type Ig domain whereas the others six domains proximal to the membrane are C2-type Ig domains (2). The cytoplasmic domain of CD22 includes six tyrosine residues that are possible targets for phosphorylation. Some regions of the intracytoplasmic tail are homologous to the tyrosine-based activations motifs (ITAM) and some others to the tyrosine-based inhibition motifs (ITIM) (2, 3).

CD22 appears constitutively associated with the BCR (B Cell antigen Receptor) and this may involve CD22 recognition of membrane IgM carbohydrate determinants (4–6). The CD22 molecule mediates adhesion of B-B lymphocyte interactions, and B cells and erythrocytes or leucocytes interactions (2, 5, 7, 8).

The CD22 antigen is detected in the cytoplasm early during B cell ontogeny (late pro-B stage), appears on the cell surface simultaneously with the expression of membrane IgD, and is found on most mature B lymphocytes (1). The CD22 antigen is lost during the terminal stages of differentiation prior to the plasma cell stage (1). On peripheral whole blood, the expression of CD22 antigen is restricted to B lymphocytes.

The SJ10.1H11 monoclonal antibody has been assigned to the CD22 cluster of differentiation at the 2nd International Workshop on Human Leukocyte Differentiation Antigens in Boston, USA, in 1984 (9).

REAGENT

IOTest CD22-PC7 Conjugated antibody
PN A74788 - 100 tests - Liquid - 10 µL/test*

Clone	SJ10.1H11
Isotype	IgG1, Mouse
Immunogen	Human NALM-1 cell line
Hybridoma	SP2/0 x spleen B cells
Source	Ascites fluid
Purification	Protein A affinity chromatography
Conjugation	Phycoerythrin-Cyanin 7 (PC7)
Molar Ratio	PC7 / Ig : 0.5 - 1.5
Fluorescence	Excites at 488 nm Emits at 770 nm

REAGENT CONTENTS

This antibody is provided in phosphate-buffered saline, containing 0.1% sodium azide and 2 mg/mL bovine serum albumin.

STATEMENTS OF WARNING

1. This reagent contains 0.1% sodium azide. Sodium azide under acid conditions yields hydrazoic acid, an extremely toxic compound. Azide compounds should be flushed with running water while being discarded. These precautions are recommended to avoid deposits in metal piping in which explosive conditions can develop. If skin or eye contact occurs, wash excessively with water.
2. Specimens, samples and all material coming in contact with them should be handled as if capable of transmitting infection and disposed of with proper precautions.
3. Never pipet by mouth and avoid contact of samples with skin and mucous membranes.
4. Do not use antibody beyond the expiration date on the label.
5. Do not expose reagents to strong light during storage or incubation.
6. Avoid microbial contamination of reagents or incorrect results might occur.
7. Use good laboratory practices when handling this reagent.

STORAGE CONDITIONS AND STABILITY

This reagent is stable up to the expiration date when stored at 2 – 8°C. Do not freeze.

REAGENT PREPARATION

No reconstitution is necessary. This monoclonal antibody may be used directly from the vial. Bring reagent to 18 – 25°C prior to use.

PRECAUTIONS

Due to the tandem structure of the fluorochrome, PC7 also emits light at 575 nm. This secondary emission peak varies from lot-to-lot of PC7. Therefore, for multi-color analysis, the compensation matrix should be carefully checked when changing the lot of a PC7-conjugate.

SELECTED RESEARCH REFERENCES

1. Kehrl, J., "CD22 workshop Panel report", 1995, Leucocyte Typing V, White Cell Differentiation Antigens. Schlossman, S.F., et al., Eds., Oxford University Press, 523-527.
2. Tedder, T.F., Tuscano, J., Sato, S., Kehrl, J.H., "CD22, A B lymphocyte-specific adhesion molecule that regulates antigen receptor signaling", 1997, Rev. Immunol., 15, 481 -504.

3. Unkeless, J.C., Jin, J., "Inhibitory receptors, ITIM sequences and phosphatases", 1997, Curr. Opin. Immunol., 9, 338-343.
4. Buhl, A.M., Cambier, J.C., "Co-receptor and accessory regulation of B-cell antigen receptor signal transduction", 1997, Immunol. Rev., 160, 127-138.
5. Law, C.L., Sidorenko, S.P., Clark, E.A., "Regulation of lymphocyte activation by the cell-surface molecule CD22", 1994, Immunol. Today, 9, 15, 442-449.
6. Doody, G.M., Dempsey, P.W., Fearon, D.T., "Activation of B lymphocytes : integrating signals from CD19, CD22 and FcγRIIb1", 1996, Curr. Opin. Immunol., 8, 378-382.
7. Lynn Wilson, G., "Genomic structure and chromosomal mapping of the human CD22 gene", 1993, J. Immunol., 11, 150, 5013.
8. Stamenkovic, I., Sgroi, D., Aruffo, A., Sy, M.S., Anderson, T., "The B lymphocyte adhesion molecule CD22 interacts with leukocyte common antigen CD45RO on T cells and alpha2-6 sialyltransferase, CD75, on B cells", 1991, Cell, 66, 1133-1144.
9. Nadler, L.M., "B cell/Leukemia panel workshop: Summary and comments", 1986, Leucocyte Typing II, Human T lymphocytes, Reinherz, E.L., et al. Eds., 4-43.

TRADEMARKS AND PATENT

Beckman Coulter, the Beckman Coulter logo, COULTER, and IOTest are registered trademarks of Beckman Coulter, Inc.

MANUFACTURED BY :

IMMUNOTECH SAS
a Beckman Coulter Company
130, avenue de Lattre de Tassigny
B.P. 177 - 13276 Marseille Cedex 9
France

For additional information in the USA, call 800-526-7694.

Outside the USA, contact your local Beckman Coulter representative.

www.beckmancoulter.com

Printed in France.
Made in France.

©2009 Beckman Coulter, Inc.

(*): 10 µL is the quantity of product sufficient to stain
5 x 10⁵ cells in a standard immunofluorescence assay

