

Monoclonal Antibody Anti-Human IgG (Pan)

PN IM0279 – Purified – Freeze-dried – 0.2 mg – Clone 8a4

For Research Use Only. Not for use in diagnostic procedures.

SPECIFICITY

The 8a4 antibody recognizes the Fc portion of all human IgG subclasses.

REAGENT

Monoclonal Antibody Anti-Human IgG (Pan)
PN IM0279 – Purified – Freeze-dried – 0.2 mg

Clone 8a4

Isotype IgG1 κ , mouse

Immunogen Purified human polyclonal IgG

Hybridoma P3-NS1/Ag1 x Balb/c spleen cells

Source Ascites fluid

Purification Ion exchange or affinity chromatography

Buffer 1 mg/mL bovine serum albumin in phosphate-buffered saline

APPLICATION

Detection of surface and intracytoplasmic immunoglobulins of human B cells by indirect staining.

Quantification of IgG by nephelometry, turbidimetry and gel precipitation techniques.

STATEMENT OF WARNINGS

1. Specimens, samples and all material coming in contact with them should be handled as if capable of transmitting infection and disposed of with proper precautions.

2. Never pipet by mouth and avoid contact of samples with skin and mucous membranes.
3. Do not use antibody beyond the expiration date on the label.
4. Avoid microbial contamination of reagents or incorrect results might occur.
5. Use good laboratory practices when handling this reagent.

STORAGE CONDITIONS AND STABILITY

This freeze-dried form may be stored at 2 – 8°C until the expiration date stated on the vial label.

No preservative has been added.

REAGENT PREPARATION

Depending of usage, reconstitute with 1 mL of distilled water, with or without 0.1% sodium azide (w/v).

The reconstituted form including 0.1% sodium azide may be stored for up to one month at 2 – 8°C.

The reconstituted form without sodium azide can be stored at –20°C or less, until the expiration date stated on the vial label.

In this case, aliquotting is recommended to avoid multiple freezing / thawing cycles.

PROCEDURE

For each application, it is recommended to establish the right range of antibody dilutions to be used for the experiment.

SELECTED RESEARCH

REFERENCES

1. Lowe, J., Bind, P., Hardie, D., Jefferis, R., Ling, N.R., "Monoclonal antibodies determinants on human gamma chains: properties of antibodies showing subclass restriction or subclass specificity", 1982, Immunology., 47, 329-336.

PRODUCT AVAILABILITY

Monoclonal Antibody Anti-Human IgG (Pan)
PN IM0279 – Purified – Freeze-dried – 0.2 mg

For additional information in the USA, call 800-526-7694.

Outside the USA, contact your local Beckman Coulter representative.

www.beckmancoulter.com

TRADEMARKS

Beckman Coulter and the Beckman Coulter logo are trademarks of Beckman Coulter, Inc.

Manufactured by:
Immunotech, a Beckman Coulter Company
130, avenue de Lattre de Tassigny, B.P. 177
13276 Marseille Cedex 9, France

©2005 Beckman Coulter, Inc.
All Rights Reserved