

PN IM0834

**Goat F(ab')₂ Fragment
Anti-Rabbit IgG (H+L)-TRITC**

Lot Number: 54546.1

For Research Use Only. Not For Use In Diagnostic Procedures.

PRODUCT

Rhodamine (TRITC)-conjugated AffiniPure F(ab')₂ Fragment Goat Anti-Rabbit IgG (H+L) (minimal cross-reaction to Human Serum Proteins)

ANTIBODY SPECIFICITY

Based on immunoelectrophoresis, the antibody reacts with the heavy chains on rabbit IgG and with light chains common to most rabbit immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. This antibody has been tested by ELISA and/or solid-phase adsorbed to ensure minimal cross-reaction with human serum proteins, but the antibody may cross-react with immunoglobulins from other species.

PURITY

The antibody was isolated from antisera by a combination of pepsin digestion and immunoaffinity chromatography using antigens coupled to agarose beads. Fc fragments and whole IgG molecules have been removed.

FORMAT

Freeze-dried powder

SIZE

1.0 mg (This product has been overfilled to ensure total recovery of stated quantity.)

PRODUCT CONCENTRATION

1.5 mg/mL

BUFFER AND STABILIZER

0.01 M Sodium Phosphate, 0.25 M NaCl, pH 7.6, containing 15 mg/mL Bovine Serum Albumin

PRESERVATIVE0.05% Sodium Azide (NaN₃)

(Note: Due to tightened regulations in many countries and states, thimerosal, a mercury-containing preservative, will no longer be included in our products. Please follow our recommendations for storage to avoid microbial contamination.)

FLUOROCHROME

Type: Tetramethyl Rhodamine Isothiocyanate (TRITC)

Fluorochrome/Protein absorbance Ratio: A550/A280 = 0.57

Absorption maximum: 550 nm

Emission maximum: 570 nm

RECONSTITUTION AND STORAGE

Store freeze-dried product at 2–8°C until reconstitution. When ready to use, reconstitute with 0.75 mL distilled water.

Centrifuge product if the solution is not clear after reconstitution. To judge clarity, draw product into a clean Pasteur pipette. Product is stable for about 6 weeks at 2–8°C as an undiluted liquid. Storing the product for more than one day at final working dilutions is not recommended. For extended storage as a liquid, add an equal volume of glycerol (ACS or better grade) for a final concentration of 50% followed by storage at –20°C. Please note that the concentration of protein and buffer salts will be one-half of the original after the addition of glycerol.

SUGGESTED DILUTION RANGE

1:50 – 1:200 for most applications

EXPIRATION DATE

One year from date of reconstitution

STATEMENT OF WARNINGS

1. For research use only. Not for use in diagnostic procedures.
2. This reagent contains 0.05% sodium azide. Sodium azide under acidic conditions yields hydrazoic acid, an extremely toxic compound. Azide compounds should be flushed with running water while being discarded. These precautions are recommended to avoid deposits in metal piping in which explosive conditions can develop. If skin or eye contact occurs, wash excessively with water.
3. Specimens, samples and all material coming in contact with them should be handled as if capable of transmitting infection and disposed of with proper precautions.
4. Never pipet by mouth and avoid contact of samples with skin and mucous membranes.
5. Do not use this reagent beyond the expiration date and after dilution no more than one day.
6. Do not expose this reagent to strong light during storage or incubation.
7. Avoid microbial contamination of reagents or incorrect results might occur.

MANUFACTURED FOR:

Immunotech, a Beckman Coulter Company
130, avenue de Lattre de Tassigny - B.P. 177
13276 Marseille Cedex 9, France
Tel: (33) +4 91 17 27 00
Fax: (33) +4 91 17 43 58
<http://www.beckmancoulter.com>