

MONOCLONAL ANTIBODY

CD61

Cat. No.	Form	Quantity	Presentation
2116	Pre-diluted	6 mL	Ready-to-use

Clone	SZ21
Isotype	IgG1 (mouse)
Immunogen	Washed human platelets (1).
Reagent preparation	Pre-diluted antibody solution is ready-to-use and should be employed according to experimental conditions and procedures validated by each individual laboratory.
Purity	Purified Ig in 50 mM Tris-HCl, 150 mM NaCl, pH 7.2 with 1 mg/mL bovine serum albumin and 0.1% sodium azide. The buffer contains a green dye.
Storage	2-8°C until the expiration date stated on the vial label.
Warning	Liquid reagents contain 0.1% sodium azide. Under acidic conditions, sodium azide yields hydrazoic acid, an extremely toxic compound. Azide compounds should be flushed with copious amounts of water while being discarded in the drain. These precautions are recommended to avoid deposits in metal piping in which explosive conditions can develop.
Biological specimens	Cytospins, smears and sections of frozen or fixed, paraffin-embedded tissues.
References	1) Ruan, C., Du, X., Hwan, Hu, X., Xi, X. & Li, P., "Characterization of the fibrinogen binding sites using monoclonal antibodies to human platelet membrane glycoproteins IIb/IIIa" 1987, <u>Thrombosis and Hemostasis</u> , 58 , 1, 243 (abstract).

December 20, 1996

MA001

FOR RESEARCH USE ONLY - NOT FOR USE IN DIAGNOSTIC PROCEDURES

**IMMUNOTECH**
A COULTER COMPANYB.P. 177 - 13276 Marseille Cedex 9 France
Tel. (33) 91 17 27 00 - Fax. (33) 91 41 43 58