

IOtest CD2-APC-Alexa Fluor 750

PN B01681 – 0.5 mL – Liquid – 10 µL/test – Clone 39C1.5

Analyte Specific Reagent.

Analytical and performance characteristics are not established.

SPECIFICITY

The CD2 antigen (1) has been formerly described as the sheep E-rosette receptor and is alternatively known as T11 antigen or leucocyte function-associated molecule 2 (LFA-2). It is a 50 kDa single chain type I transmembrane glycoprotein that comprises two external domains belonging to the immunoglobulin superfamily (IgSF) (2). Crystallographic studies of soluble, deglycosylated forms of rat and human CD2 molecules (3, 4), revealed that the N-terminal, distal domain 1 of the molecule is a V-type IgSF domain, and that the proximal domain 2 is a C-type IgSF domain. The V-type domain 1 lacks the usually conserved disulphide bonds between the beta sheets. The extracellular segment includes at least three N-glycosylation sites, believed to be involved in the interactions with the CD2 ligands. CD2 has a relatively large cytoplasmic domain which is required for the activation produced by certain combinations of CD2 antibodies. Signaling molecules, such as Fyn, Lck and the PI3-kinase, have been reported to associate with the intracellular domain of CD2 (5).

The first CD2 ligand identified was CD58 (LFA-3), a heavily glycosylated molecule of 70 kDa, broadly expressed on leucocytes, erythrocytes and endothelial cells (6). The binding site for CD58 lies on one face of the first domain of human CD2 (7). In addition to CD58, there is evidence that CD48 and CD59 may be ligands for CD2. The CD2 molecule is also able to interact with CD48, but with considerably weaker affinity than with CD58 (8).

The CD58-CD2 interaction is a well known component of intercellular adhesion and costimulatory signaling in T cells (6, 9). In contrast, little is known about the physiological roles of the other putative ligands (1). CD2 has been described as being involved in the regulation of human T-cell cytokine production (10), and in a Fas-independent induced apoptosis of activated human peripheral T cells (11 – 13). CD2 is present on all human non-B peripheral lymphocytes, on the majority of thymic T cells (14, 15), and on a subset of thymic B cells.

Several epitopes can be distinguished on the CD2 molecule (16). The 39C1.5 monoclonal antibody, also known as CD2.9 (17), reacts with the T11-1 group of epitopes.

It has been assigned to the CD2 cluster of differentiation at the 2nd International Workshop on Human Leucocyte Differentiation Antigens in Boston, USA, in 1984 (16, 18).

REAGENT

IOtest CD2-APC-Alexa Fluor 750
Conjugated antibody
PN B01681 - 0.5 mL - Liquid - 10 µL/test

Clone	39C1.5
Isotype	IgG2a, Rat
Immunogen	PHA stimulated human lymphoblasts
Hybridoma	X63 x rat
Source	Ascites fluid or supernatant of in vitro cultured hybridoma cells.
Purification	Ion exchange or affinity chromatography
Conjugation	Allophycocyanin-Alexa Fluor 750 (APC-Alexa Fluor)
Molar Ratio	APC-Alexa Fluor / Ig : 0.5 - 1.5
Fluorescence	Excites at 633/638 nm Emits at 775 nm

REAGENT CONTENTS

This antibody is provided in phosphate-buffered saline, containing 0.1% sodium azide and 2 mg/mL bovine serum albumin.

STATEMENTS OF WARNING

1. This reagent contains 0.1% sodium azide. Sodium azide under acid conditions yields hydrazoic acid, an extremely toxic compound. Azide compounds should be flushed with running water while being discarded. These precautions are recommended to avoid deposits in metal piping in which explosive conditions can develop. If skin or eye contact occurs, wash excessively with water.
2. Specimens, samples and all material coming in contact with them should be considered potentially infectious and disposed of with proper precautions.
3. Never pipet by mouth and avoid contact of samples with skin and mucous membranes.
4. Do not use antibody beyond the expiration date on the label.
5. Do not expose reagents to strong light during storage or incubation.
6. Avoid microbial contamination of reagents or incorrect results might occur.
7. Use good laboratory practices when handling this reagent.

STORAGE AND HANDLING CONDITIONS AND STABILITY

This reagent is stable up to the expiration date when stored at 2 – 8°C. Do not freeze.

No reconstitution is necessary. This monoclonal antibody may be used directly from the vial. Bring reagent to 18 – 25°C prior to use.

PRECAUTIONS

Due to the tandem structure of the fluorochrome, APC-Alexa Fluor 750 also emits light at 660 nm. This secondary emission peak varies from lot-to-lot of APC-Alexa Fluor 750. Therefore, for multi-color analysis, the compensation matrix should be carefully checked when changing the lot of a APC-Alexa Fluor 750-conjugate.

SELECTED RESEARCH REFERENCES

1. Davis, S.J., van Der Merwe, P.A., "The structure and ligand interactions of CD2 : implications for T-cell function", 1996, *Immunol. Today*, 17, 177-187.
2. Sewel, W.A., Brown, M.H., Dunne, J., Owen, M.J., Crumpton, M.J., "Molecular cloning of the human T lymphocyte surface CD2 (T11) antigen", 1986, *Proc.Natl. Acad.Sci. USA*, 83, 8718-8722.
3. Jones, E.Y., Davis, S.J., Williams, A.F., Harlos, K., Stuart, D.I., "Crystal structure at 2.8 Å resolution of a soluble form of the cell adhesion molecule CD2", 1992, *Nature*, 360, 232-239.
4. Bodian, D.L., Jones, E.Y., Harlos, K., Stuart, D.I., Davis, S.J., "Crystal structure of the extracellular region of the human cell adhesion molecule CD2 at 2.5 Å resolution", 1994, *Structure*, 2, 755-766.
5. Thèze, J., Alzari, P.M., Bertoglio, J., "Interleukin 2 and its receptors: recent advances and new immunological functions", 1996, *Immunol. Today*, 17,481-486.
6. Dustin, M.L., Sanders, M.E., Shaw, S., Springer, T.A., "Purified lymphocyte function-associated antigen 3 binds to CD2 and mediates T lymphocyte adhesion", 1987, *J. Exp. Med.*, 165, 677-392.
7. Somoza, C., Driscoll, P.C., Cyster, J.G., Williams, A.F., "Mutational analysis of the CD2/CD58 interaction : the binding site for CD58 lies on one face of the first domain of human CD2", 1993, *J. Exp. Med.*, 178, 549-558.
8. Arulanandam, A.R., Moingeon, P., Concino, M.F., Recny, M.A., Kato, K., Yagita, H., Koyasu, S., Reinherz, E.L., "A soluble multimetric recombinant CD2 protein identifies CD48 as a low affinity ligand for human CD2 : divergence of CD2 ligands during the evolution of humans and mice", 1993, *J. Exp. Med.*, 177, 1439-1450.

9. Denning, S.M., Dustin, M.L., Springer, T.A., Singer, K.H., Haynes, B.F., "Purified lymphocyte function-associated antigen-3 (LFA-3) activates human thymocytes via the CD2 pathway", 1988, *J. Immunol.*, 141, 2980-2985.
10. Holter, W., Schwarz, M., Cerwenka, A., Knapp, W., "The role of CD2 as a regulator of human T-cell cytokine production", 1996, *Immunol. Rev.*, 153, 107-122.
11. Rouleau, M., Bernard, A., Lantz, O., Vernant, J.P., Charpentier, B., Senik, A., "Apoptosis of activated CD8+/CD57+ T cells is induced by some combinations of anti-CD2 mAb", 1993, *J. Immunol.*, 151, 3547-3556.
12. Mollereau, B., Deckert, M., Déas, O., Rieux-Laucat, F., Hirsch, F., Bernard, A., Fischer, A., Lynch, D.H., Charpentier, B., Le Deist, F., Senik, A., "CD2-induced apoptosis in activated human peripheral T cells", 1996, *J. Immunol.*, 156, 3184-3190.
13. Rouleau, M., Mollereau, B., Bernard, A., Metivier, D., Rosenthal-Allieri, M.A., Charpentier, B., Senik, A., "CD2 induced apoptosis of peripheral T cells", 1997, *Transplant. Proc.*, 29, 2377-2378.
14. Bernard, A., Brottier, P., Georget, E., Lepage, V., Boumsell, L., "The epitopic dissection of the CD2 defined molecule: relationship of the second workshop antibodies in terms of reactivities with leukocytes, rosette blocking properties, induction of positive modulation of the molecule, and triggering T cell activation", 1985, *Leucocyte Typing II, Human T lymphocytes*, Reinherz, E.L., et al. Eds., 55-59.
15. Bierber, B.E., Bogart, R.E., Wolff, I.H., Burkakoff, S.J., "Functional analysis of CD2 Mab reactivity", 1989, *Leucocyte Typing IV, White Cell Differentiation Antigens*. W. Knapp, et al., Eds., Oxford University Press, 274-277.
16. Haynes, B.F., "Summary of T cell studies", 1986, in *Leucocyte Typing II, White Cell Differentiation Antigen*, Rheinherz, E.L., et al., Eds., Springer-Verlag, p. 3-30.
17. Olive, D., Ragueneau, M., Cerdan, C., Dubreuil, P., Lopez, M., Mawas, C., "Anti-CD2 (Sheep red blood cell receptor) monoclonal antibodies and T-cell activation. Pairs of anti-T11-1 and T11-2 (CD2 subgroups) are strongly mitogenic for T cells in presence of 12-0 tetradecanoylphorbol 13 acetate", 1986, *Eur. J. Immunol.*, 16, 1063-1068.
18. Knowles R. W., "Immunochemical Analysis of the T cell-specific antigens", 1986, in *Leucocyte Typing II, White Cell Differentiation Antigen*, Rheinherz, E.L., et al., Eds., Springer-Verlag, p. 266-271.

MANUFACTURED BY :

IMMUNOTECH SAS
a Beckman Coulter Company
130, avenue de Lattre de Tassigny
B.P. 177 - 13276 Marseille Cedex 9
France

For additional information in the USA, call
800-526-7694.
Outside the USA, contact your local Beckman
Coulter representative.

www.beckmancoulter.com

Printed in France.
Made in France.

©2011 Beckman Coulter, Inc.

TRADEMARKS

Beckman Coulter logo, ECD, and IOTest, are trademarks of Beckman Coulter; Beckman Coulter logo, IOTest are registered in the USPTO and SIPO.

Alexa Fluor is a trademark of Molecular Probes, Inc.