

IOTest CD19-Alexa Fluor 700

PN B76283 – 0.5 mL – Liquid – Clone J3.119

Analyte Specific Reagent.

Analytical and performance characteristics are not established

IOTest

CD19-Alexa Fluor 700

PN B76283

Conjugated Antibody
Liquid - 0.5 mL

Specifications	
Clone	J3.119
Hybridoma	NS1 x balb/c
Immunogen	SKLY 18 lymphoma cells
Isotype	IgG1 kappa
Species	Mouse
Source	Ascites fluid or supernatant of in vitro cultured hybridoma cells.
Purification	Affinity chromatography
Fluorochrome	Alexa Fluor 700
Molar ratio	Alexa Fluor 700 / Ig: 3 - 5.5
λ excitation	695 nm
Emission Peak	720 nm
Buffer	PBS pH 7.2 plus 2 mg / mL BSA and 0.1% NaN ₃

SPECIFICITY

The CD19 antigen (also called B4) is a type I membrane glycoprotein with a molecular weight of 95 kDa (1, 2).

Its extracellular domain comprises 280 amino acids organized as two C2-type Ig-like domains separated by a smaller potentially disulfide-linked domain. The extensive cytoplasmic domain of CD19 contains nine conserved tyrosine residues, and several of these are located within potential src-homology region 2 (SH2)-binding sites.

CD19 is a signal transduction molecule that regulates lymphocyte development, activation, and differentiation (3, 4).

The molecule is expressed on all normal B lymphocytes including pro-B lymphocytes, but it is lost in maturation to plasma cells (3, 5).

It is also found on the surface of follicular dendritic cells, on the early cells of myelomonocytic lineage and on most stabilized B cell lines. It is not present on normal T lymphocytes, NK cells, monocytes, and granulocytes.

CD19 can be associated within the membrane to form hetero-oligomeric structures with other surface molecules including CD21, the complement receptor type 2 (CR2), and CD81 (TAPA-1) (6). The extracellular and transmembrane domains of CD19 are required for the interaction of this molecule with CD21 and CD81. Co-ligation of the CD19-CD21-CD81 complex with the surface IgM-B-cell antigen receptor (BCR) leads to the phosphorylation of CD19 by Syk (6, 7), followed by the recruitment, through tyrosine phosphorylated CD19, of positive signal transduction effectors such as phosphatidylinositol 3 kinase (PI3 kinase), Lyn, and Fyn (8, 9).

In vitro studies show that the CD19 antibodies have an inhibitory effect on the activation and proliferation of B lymphocytes. They also inhibit the B cell response after

co-stimulation by anti-immunoglobulin and interleukin 4.

The J3-119 monoclonal antibody (mAb) reacts with B lymphocytes, not with T lymphocytes, monocytes or granulocytes. It stains mantle-zone and germinal-center lymphoid cells, as well as follicular dendritic cells.

Plasma cells are negative.

The J3-119 mAb has been assigned to the CD19 cluster of differentiation during the fourth International Workshop on Human Leucocyte Differentiation Antigens held in Vienna, Austria in 1989 (1, 2).

REAGENT CONTENTS

Concentration: See lot specific Certificate of Analysis at www.beckmancoulter.com.

PRECAUTIONS

1. This reagent contains 0.1% sodium azide. Sodium azide under acid conditions yields hydrazoic acid, an extremely toxic compound. Azide compounds should be flushed with running water while being discarded. These precautions are recommended to avoid deposits in metal piping in which explosive conditions can develop. If skin or eye contact occurs, wash excessively with water.
2. Specimens, samples and all material coming in contact with them should be considered potentially infectious and disposed of with proper precautions.
3. Never pipet by mouth and avoid contact of samples with skin and mucous membranes.
4. Do not use antibody beyond the expiration date on the label.
5. Do not expose reagents to strong light during storage or incubation.
6. Avoid microbial contamination of reagents or incorrect results might occur.

7. Use good laboratory practices when handling this reagent.
8. Any change in the physical appearance of the reagents may indicate deterioration and the reagent should not be used.

STORAGE AND HANDLING CONDITIONS AND STABILITY

This reagent is stable up to the expiration date when stored at 2 – 8°C. Do not freeze. No reconstitution is necessary. This monoclonal antibody may be used directly from the vial. Bring reagent to 18 – 25°C prior to use.

SELECTED RESEARCH REFERENCES

1. "CD Guide " Compiled by the organizing committee, 1989, Leucocyte Typing IV, White Cell Differentiation Antigens. W. Knapp, et al., Eds., Oxford University Press, 1078.
2. "Listing of all Fourth Workshop antibodies", 1989, Leucocyte Typing IV, White Cell Differentiation Antigens. W. Knapp, et al., Eds., Oxford University Press, 1094-1110.
3. Doody, G.M., Dempsey, P.W., Fearon, D.T., "Activation of B lymphocytes : integrating signals from CD19, CD22 and Fc γ RIIb1", 1996, Cur. Opin. Immunol., 8, 378-382.
4. Pesando, J. M., Bouchard, L. S., McMaster, B. E., "CD19 is functionally and physically associated with surface immunoglobulin", 1989, J. Exp. Med., 170, 2159-2164.
5. Loken, M.R., Shah, V.O., Dattilio, K.L., Civin, C.I., "Flow cytometric analysis of human bone marrow. II. Normal B lymphocyte development", 1987, Blood, 70, 1316-1324.
6. Bradbuys, L.E., Kansas, G.S., Levy, S., Evans, R.L., Tedder, T.F., "The CD19/CD21 signal transducing complex of human B lymphocytes includes the

IOTest CD19-Alexa Fluor 700

PN B76283 – 0.5 mL – Liquid – Clone J3.119

target of antiproliferative antibody-1 and Leu-13 molecules", 1992, J. Immunol., 149, 2841-2850.

7. Carter, R.H., Doody, G.M., Bolen, J.B., Fearon, D.T., "Membrane IgM-induced tyrosine phosphorylation of CD19 requires a CD19 domain that mediates association with components of the B cell antigen receptor complex", 1997, J. Immunol., 158, 3062-3069.
8. Tedder, T.F., Zhou, L.J., Engel, P., "The CD19/CD21 signal transduction complex of B lymphocytes", 1994, Immunol. Today, 15, 437-442.
9. Kurosaki, T., "Molecular mechanisms in B cell antigen receptor signaling", 1997, Curr. Opin. Immunol., 9, 309-318

TRADEMARKS

Beckman Coulter, the stylized logo and IOTest, are trademarks of Beckman Coulter, Inc., and registered with the USPTO.

Alexa Fluor is a trademark of Molecular Probes, Inc.

MANUFACTURED BY:

IMMUNOTECH SAS
a Beckman Coulter Company
130, avenue de Lattre de Tassigny
B.P. 177 - 13276 Marseille Cedex 9
France

For additional information, or if damaged product is received, call Beckman Coulter Customer Service at 800-526-7694 (USA or Canada) or contact your local Beckman Coulter Representative.

www.beckmancoulter.com

Printed in France.
Made in France.

©2015 Beckman Coulter, Inc.
All Rights Reserved